



Official reprint from UpToDate®

www.uptodate.com ©2018 UpToDate, Inc. and/or its affiliates. All Rights Reserved.

The content on the UpToDate website is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your own physician or other qualified health care professional regarding any medical questions or conditions. The use of UpToDate content is governed by the [UpToDate Terms of Use](#). ©2018 UpToDate, Inc. All rights reserved.

Lithium: Patient drug information

Copyright 1978-2018 Lexicomp, Inc. All rights reserved.

(For additional information [see "Lithium: Drug information"](#) and [see "Lithium: Pediatric drug information"](#))

Brand Names: US Lithobid

Brand Names: Canada Carbolith; Lithane; Lithmax

Warning

- This drug may cause bad side effects at doses that are close to the normal dose. Do not take more of this drug than you are told. You will have blood tests to check the level of this drug in your blood. Tell your doctor right away if you feel very sleepy, pass more urine than normal, or have a heartbeat that does not feel normal, loose stools (diarrhea), throwing up, blurred eyesight, ringing in the ears, giddiness, weak muscles, shakiness, twitching, seizures, eye movements that are not normal, change in balance, or very upset stomach.

What is this drug used for?

- It is used to treat bipolar problems.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you have an allergy to lithium or any other part of this drug.
- If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If you have any of these health problems: Fluid loss (dehydrated), heart disease, kidney disease, low levels of salt in your blood, or if you are very sick or weak.
- If you have a certain heart problem called Brugada syndrome or you have ever passed out for an unknown reason.

- If a family member has Brugada Syndrome or has suddenly died before 45 years of age.
- If you are taking drugs for blood pressure or heart problems. You may need to have your blood work checked more closely.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- If you start on a low-salt diet, tell your doctor. It will change the way your body handles the lithium.
- Talk with your doctor before you drink alcohol.
- Fever, infection, throwing up, loose stools (diarrhea), or sweating a lot may change how much of this drug is in your blood. If any of these happen, talk with the doctor.
- Be careful in hot weather or while being active. Drink lots of fluids to stop fluid loss.
- Tell your doctor if you have signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- A certain heart problem (Brugada Syndrome) has shown up in some people taking this drug. These people had this problem but did not know it at the time. People with Brugada Syndrome have a heartbeat that is not normal and are at risk for sudden death. Talk with the doctor.
- Brain problems have rarely happened in people taking this drug with certain other drugs like haloperidol. Sometimes, these brain problems have led to long-lasting brain damage. Talk with the doctor.
- A very bad and sometimes deadly health problem called serotonin syndrome may happen. The risk may be greater if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea, upset stomach, or throwing up; or very bad headache.
- Raised pressure in the brain and swelling in the eye has happened with this drug. This can lead to eyesight problems like loss of eyesight. Call your doctor right away if you have a very bad headache; ringing in the ears; or eyesight problems like blurred eyesight, seeing double, or loss of eyesight.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug is not approved for use in children younger than 12 years old. Talk with the doctor.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding or plan to breast-feed. This drug passes into breast milk and may harm your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of too much acid in the blood (acidosis) like confusion; fast breathing; fast heartbeat; a heartbeat that does not feel normal; very bad stomach pain, upset stomach, or throwing up; feeling very sleepy; shortness of breath; or feeling very tired or weak.
- Signs of thyroid problems like a change in weight without trying, feeling nervous and excitable, feeling restless, feeling very weak, hair thinning, low mood (depression), neck swelling, not able to focus, not able to handle heat or cold, period (menstrual) changes, shakiness, or sweating.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat, more thirst, seizures, feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Dizziness or passing out.
- Slow heartbeat.
- Feeling confused.
- Fever.
- Trouble controlling body movements.
- Hallucinations (seeing or hearing things that are not there).
- Memory problems or loss.
- Restlessness.
- Trouble walking.
- Swelling in the feet or hands.
- A change in weight without trying.
- Slurred speech.
- Not able to control bladder.
- Not able to control bowels.
- Sex problems.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Shakiness of the hand.

- Feeling tired or weak.
- Upset stomach.
- Feeling sleepy.
- Not hungry.
- Belly pain.
- More saliva.
- Gas.
- Dry mouth.
- Headache.
- Dry or thin hair.
- Hair loss.
- Change in taste.
- Joint pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

All products:

- Take with or without food. Take with food if it causes an upset stomach.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.
- Keep your salt use the same from day to day.

Liquid (solution):

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

Long-acting products:

- Swallow whole. Do not chew or crush.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- If you are not sure what to do if you miss a dose, call your doctor.

How do I store and/or throw out this drug?

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor.
- Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Use of UpToDate is subject to the [Subscription and License Agreement](#).

Topic 10813 Version 141.0