



The content on the UpToDate website is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your own physician or other qualified health care professional regarding any medical questions or conditions. The use of UpToDate content is governed by the [UpToDate Terms of Use](#). ©2018 UpToDate, Inc. All rights reserved.

Paroxetine: Patient drug information

Copyright 1978-2018 Lexicomp, Inc. All rights reserved.

(For additional information [see "Paroxetine: Drug information"](#) and [see "Paroxetine: Pediatric drug information"](#))

Brand Names: US Brisdelle; Paxil; Paxil CR; Pexeva

Brand Names: Canada Paxil; Paxil CR

Warning

- Drugs like this one have raised the chance of suicidal thoughts or actions in children and young adults. The risk may be greater in people who have had these thoughts or actions in the past. All people who take this drug need to be watched closely. Call the doctor right away if signs like low mood (depression), nervousness, restlessness, grouchiness, panic attacks, or changes in mood or actions are new or worse. Call the doctor right away if any thoughts or actions of suicide occur.
- This drug is not approved for use in children. Talk with the doctor.

What is this drug used for?

- It is used to treat low mood (depression).
- It is used to treat obsessive-compulsive problems.
- It is used to treat panic attacks.
- It is used to treat social anxiety problems.
- It is used to treat anxiety.
- It is used to treat post-traumatic stress.
- It is used to treat mood problems caused by monthly periods.
- It is used to treat hot flashes caused by menopause.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

For all uses of this drug:

- If you have an allergy to paroxetine or any other part of this drug.
- If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If you are taking any of these drugs: Linezolid, methylene blue, pimozone, or thioridazine.
- If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure.

For treating hot flashes caused by menopause:

- If you are pregnant or may be pregnant. Do not take this drug if you are pregnant.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert or have clear eyesight until you see how this drug affects you.
- Do not stop taking this drug all of a sudden without calling your doctor. You may have a greater risk of side effects. If you need to stop this drug, you will want to slowly stop it as ordered by your doctor.
- Avoid drinking alcohol while taking this drug.
- Talk with your doctor before you use other drugs and natural products that slow your actions.
- It may take several weeks to see the full effects.
- This drug may affect how much of some other drugs are in your body. If you are taking other drugs, talk with your doctor. You may need to have your blood work checked more closely while taking this drug with your other drugs.
- This drug may raise the chance of a broken bone. Talk with the doctor.
- This drug may raise the chance of bleeding. Sometimes, bleeding can be life-threatening. Talk with the doctor.
- Some people may have a higher chance of eye problems with this drug. Your doctor may want you to have an eye exam to see if you have a higher chance of these eye problems. Call your doctor right away if you have eye pain, change in eyesight, or swelling or redness in or around the eye.
- A very bad and sometimes deadly health problem called serotonin syndrome may happen. The risk may be greater if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea, upset stomach, or throwing up; or very bad headache.

- This drug can cause low sodium levels. Very low sodium levels can be life-threatening, leading to seizures, passing out, trouble breathing, or death. Talk with the doctor.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.
- This drug may affect sperm in men. This may affect being able to father a child. Talk with the doctor.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of low sodium levels like headache, trouble focusing, memory problems, feeling confused, weakness, seizures, or change in balance.
- Signs of bleeding like throwing up blood or throw up that looks like coffee grounds; coughing up blood; blood in the urine; black, red, or tarry stools; bleeding from the gums; vaginal bleeding that is not normal; bruises without a reason or that get bigger; or any bleeding that is very bad or that you cannot stop.
- Feeling confused.
- Change in balance.
- Agitation, twitching, sweating, or muscle stiffness.
- Very nervous and excitable.
- A heartbeat that does not feel normal.
- Hallucinations (seeing or hearing things that are not there).
- Very bad dizziness or passing out.
- Bone pain.
- Seizures.
- A big weight loss.
- Restlessness.
- A burning, numbness, or tingling feeling that is not normal.
- Sex problems like lowered interest in sex or ejaculation problems.
- Call your doctor right away if you have a painful erection (hard penis) or an erection that lasts for longer than 4 hours. This may happen even when you are not having sex. If this is not treated right away, it may lead to lasting sex problems and you may not be able to have sex.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Dizziness.
- Feeling sleepy.
- Feeling nervous and excitable.
- Headache.
- Upset stomach or throwing up.
- Constipation.
- Loose stools (diarrhea).
- Dry mouth.
- Not able to sleep.
- Feeling tired or weak.
- Not hungry.
- Shakiness.
- Yawning.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

All products:

- To gain the most benefit, do not miss doses.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Take with or without food.

Liquid (suspension):

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

Tablet:

- Swallow whole. Do not chew, break, or crush.

Capsule:

- Take this drug at bedtime.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.

- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Protect from light.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor.
- Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Use of UpToDate is subject to the [Subscription and License Agreement](#).

Topic 10950 Version 155.0